



## RESEARCH ARTICLE

# COVID 19 in Pregnancy: Immunology Savior or Culprit

Dr Brinderjeet Kaur\*

<sup>1</sup>Consultant Department of Obstetrics and Gynecology Santokba Durlabhji Memorial Hospital and Research Center, Jaipur, India

### Abstract

Corona pandemic has been a nightmare for health care, from the point of view of transmission, pathogenesis, diagnosis and treatment, everything in doldrums. Pregnancy in itself encompasses altered physiology, immunity and often is characterized by unpredicted bodily responses. The short review is an attempt to summarize the knowledge gained so far in context with COVID 19 infection in pregnancy. The paper highlights gaps in our present understanding and emphasize on more research for understanding the double edged sword - immune system and its response to COVID 19 infection in pregnant woman.

Keywords: Immunity, Pregnancy and COVID -19

Copyright : © 2021 The Authors. Published by Publisher. This is an open access article under the CC BY-NC-ND license (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1 | INTRODUCTION

The information pertaining to pregnancy and COVID-19 infection remains limited and controversial. COVID -19 has been proclaimed as a global pandemic respiratory disease caused by severe acute respiratory syndrome corona virus 2 (SARS-CoV-2) (1). The viral infection has been reported to have three stages. Stage 1 – The incubation period where in most of the cases are asymptomatic however the virus survives in the host undetected but infective. The incubation period from the day is usually of 5-7 days. Stage II – The virus is detectable with minor or mild symptoms like fever, cough, sore throat, myalgia, and fatigue (2–5). Stage III This is the full blown disease with cytokine storm manifesting with symptoms like high fever with coughing, headache

, difficulty in breathing, pneumonia, diarrhea, haemoptysis and excessive sputum. Stage III can be divided into severe stage- defined as tachypnoea >30 breaths per minute, oxygen saturation  $\leq 93\%$  at rest, or PaO<sub>2</sub>/ FiO<sub>2</sub> ratio < 300 mmHg and critical stage characterized by- respiratory failure requiring mechanical ventilation, septic shock or other organ dysfunction (6). This critical stage requires intensive care. The current understanding of COVID-19 pathogenesis and clinical spectrum points

**Supplementary information** The online version of this article (<https://doi.org/xx.xxx/xxx.xx>) contains supplementary material, which is available to authorized users.

**Corresponding Author:** Dr Brinderjeet Kaur  
Dr Brinderjeet Kaur

out that individuals >65 years, with compromised immune system, other underlying chronic disease like diabetes mellitus or chronic infections and perhaps pregnant comprise the vulnerable group. (7)

## 2 | IMMUNE RESPONSE IN NORMAL PREGNANCY

Pregnancy has been described as physiologically altered inflammatory state (8, 9). The immune system behaves like a pendulum between proinflammatory to anti-inflammatory state during the various stages of pregnancy. The first trimester is pro-inflammatory and second trimester been anti-inflammatory and third trimester shifts back into pro-inflammatory state. (9) An inflammatory response is triggered for the blastocyst to successfully penetrate the uterus for implantation and placentation during first trimester (9). At this stage the maternal system weakens and mother suffers from morning sickness, weakness, fatigue, headache indicating pro-inflammatory phase. (10) In the second trimester maternal health improves and anti-inflammatory state leading to fetal development and growth swiftly. In the third trimester, the development of fetus completes and immune response is again activated for the delivery of the baby and the placenta (11), as the immune cells invade the myometrium with the pro-inflammatory phase facilitates contraction in the uterus. (12)

It is important to take into consideration the role of NK cells. These NK cells protect the body from diseases by secreting cytokines such as INF  $\gamma$  and TNF $\alpha$ , which act on other immune cells like macrophages to fight the infection. NK cells are controlled by inhibitory receptors such as NKG2A. Activation of NKG2A receptors prevent NK cells from destroying trophoblast cells. (13–15) In a normal healthy pregnancy, the percentage of NK cells in the peripheral blood tend to increase in the first trimester, decrease in second and third trimester. The uterine NK cells during the first trimester become progressively less granular and decrease in number, leaving only minimal Uterine NK cells at term. Uterine NK cells are known to regulate trophoblast invasion by producing the chemokines interleukin (IL-8) and

IFN-inducible protein (IP-10) and induce vascular growth by secreting angiogenic factors (16). However, increased activation of uterine NK cells have been associated with pregnancy complications such as spontaneous abortion and pre eclampsia. (14)

## 3 | PATHOGENESIS IN COVID-19

The structure of COVID-19 has been described as a positive single stranded RNA genome having four genes; a spike protein, an envelope, a membrane and a nucleocapsid. (17) The main target of the virus is the pulmonary system where the virus binds to the host receptor, the angiotensin converting enzyme 2 (ACE 2) via receptor binding domains present on surface of epithelia cells. (15) The spike protein then undergoes a conformational modification for the viral envelope to bind to the cell membrane and release RNA into the host cells. Once RNA enters the host cell, it gets translated into viral replicases, which are split into smaller particles by enzymes called proteinases. The particles are then translated into viral proteins by mRNA and gather into virions on the endoplasmic reticulum and the Golgi apparatus where they are released out of the cells via vesicles (18). Following their release, they enter the alveoli cells, endothelial cells and blood cells. This causes exaggerated activation of immune cells and cytokines.

## 4 | CORONA VIRUS IMMUNE RESPONSE

Studies have revealed that COVID-19 is characterized by increased total neutrophils, reduced total lymphocytes, increased IL-6, increased C-reactive, D-dimer and Procalcitonin are indicate severe COVID-19 infection (4). The latter findings also correlate with disease severity and death. A study by Evangelos et al investigating the immune response revealed that patients with severe respiratory failure displayed either macrophage activation syndrome (MAS) or very low human leucocyte antigen D related (HLA-DR) expression accompanied by profound depletion of CD4 lymphocytes, CD19 lymphocytes, and natural killer (NK cells) (19).

Both NK cells and T cells (cell mediated cytotoxic adaptive immunity ) are important for control of infection , their depletion is associated with severe COVID-19 infection (20). Literature studies have noticed an increased expression of NKG2A, which is an inhibitory receptor for NK. (20, 21)

**5 | SIMILARITY IN IMMUNE RESPONSE BETWEEN CORONA INFECTION AND PREGNANCYTABLE ??**

Proinflammatory cytokines such as IL – 1, IL-6, INF  $\gamma$ , MCP 1 and IP-10 have been reported to be elevated into the blood of patients infected by COVID -19 (5). Patients with severe disease show significantly higher percentage of CD 14, CD 16 inflammatory monocytes in peripheral blood than patients with mild disease. These cells secrete inflammatory cytokines that contribute to the cytokine storm, including MCP1, IP-10 and MIP1  $\alpha$ ,TNF (5, 7). The ripple of cytokines may result in septic shock, multi organ failure which may result in myocardial damage and circulatory failure observed in some patients. (21)

The pregnancy during first and second trimester is a pro inflammatory state more so in the I trimester. Therefore the risk of contracting COVID-19 infection is higher in I trimester. Immune defense in I trimester is governed by activation of NK cells and monocytes at one end and attenuation of cell mediated Th 1 immunity with predominance of Th2 dominated environment that increases morbidity due to infection. (22)

Corona virus is known to target the host receptor ACE 2 via receptor binding domain. The expression of ACE 2 in the placenta is higher in the vilous cytotrophoblast , syncytiotrophoblast cells in the deciduas during the first trimester of the pregnancy (23). Since ACE 2 is highly expressed in this region of the placenta , this not only increases the risk of the mother contracting SARS-CoV -2 , but this is also plausible cause that transmission from the mother to child may occur. (24)Till now , no reports on mother to child transmission of the corona virus have been reported . Infact this transmission

is highly unlikely since placenta creates a protective mechanism against viruses .The placenta actually creates protection against foreign pathogens by anti-microbial action . (11) The trophoblast cells stimulate the expression of secretory leucocyte protease inhibitor (SLPI) and INF- , which are recognized for inhibitory action against viruses.

**TABLE 1:** The immune response in Normal Pregnancy vs COVID -19

Pregnancy	COVID -19 Infection
↓ Natural Killer Cells	↓ Natural Killer Cells
↑NKG2A Receptors	↑NKG2A Receptors
↓ Lymphocytes	↓ Lymphocytes
↑Pro inflammatory factors	↑Pro inflammatory factors
↑ACE 2 Receptors	↑ACE 2 Receptors

Pregnancy brings mechanical effects on physiological cardio respiratory system alterations causing diaphragmatic displacement by the gravid uterus and the gestational weight gain resulting in altered pulmonary volumes leading to reduced total lung capacity and to an inability to clear effectively the pulmonary secretions. (25) This could add to susceptibility to infection.

**6 | FACTORS LOWERING THE RISK OF INFECTION IN PREGNANCY**

Inspite of the much publicized increased risk of pregnant women, the reports of inflammatory lung process towards extensive tissue damage and the ARDS has been rarely observed. The speculated explanation could be physiological ‘silencing ‘of the Th 1 pro –inflammatory response and dominance of Th 2 over Th 1 immunity which may account for a more restrained inflammatory cascade in pregnant women with COVID19. While the Th1 lymphocyte cytokines include gamma interferon (INF –  $\gamma$ ),interleukin IL-1 $\alpha$  , IL-1, IL-12, IL-6 are microbicidal and proinflammatory . On the contrary in Th 2 dominated pregnancy state anti –inflammatory response is mediated by IL-, IL-10, IL-13 and transforming growth factor beta (TGF) (26, 27).

It is also possible that pregnancy been dominated by anti inflammatory milieu which makes the immune response less deregulated and blunt the fatal cytokine storm which would have otherwise set in by the Th 1 cells. On this ground, it may be explained why hydrochloroquine (HCQ), colchicines, monoclonal antibody anti- IL1R ( Toclimuzab and Anakinra ) have been documented to mitigate the inflammatory response and provide clinical benefits . Hydroxychloroquine, Colchicine and Toclimuzab have ability to reduce IFN- $\gamma$  and IL-1 secretion by inhibition of IL-6 receptors. (26) Heparin use is all supported due to its anti –inflammatory action which may further decrease the risk of arterial and venous thrombosis. (27–29)

## 7 | CONCLUSION

We may infer from above discussion that COVID-19 infection and pregnancy has few similar immunological responses. The I and II trimester of pregnancy are more susceptible to COVI19 disease. The current literature does not establish vertical maternal to child transmission of COVID-19 infection and therefore it can be concluded overall COVID-19 infection in pregnancy follows a benign course with less likelihood of cytokine storm or grave COVID-19 complications. Further studies on immunological aspects of COVID-19 infection in pregnant woman are required for enhancing our knowledge on this bewildering aspect of immunity , a double edged sword.

## 8 | SOURCES OF FUNDING

Nil

## 9 | CONFLICT OF INTEREST

Nil

## REFERENCES

1. Organization WH Coronavirus disease. 2020;2019:72–72.
2. Chan JF, Yuan S, Kok K, To H, Chu, Yang J. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person -to -person transmission: a study of a family cluster. *Lancet*. 2020;395:514–537.
3. Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan. China : a descriptive study. *Lancet*. 10223;2020:507–520.
4. Guan W, Liang NZY, H W, Ou CQ. HeJ x , et al Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med*;2020:1708–1728.
5. Huang C, Wang Y, Li X, Ren I, Zhaoj, Ha Y. Clinical features of patients infected with 2019 novel coronavirus in Wuhan , China. *Lancet*. 2020;395:495–506.
6. Surveillances V; 2020.
7. Rothan HA, Byrareddy SN. The epidemiology and pathogenesis of coronavirus disease (COVID-19 )outbreak. *J Autoimmun*. 2020;p. 102433–102433.
8. Wegmann TC, Lin H, Guilbert I, Mosmann TR. Bidirectional cytokine interactions in the maternal -fetal relationship : is successful pregnancy a TH2 phenomenon. *Immunol Today*. 1993;14(7):353–359.
9. Dekel N, Gnainsky Y, Granot I, Mor C. Inflammation and implantation. *Am J Reprod Immunol*. 2010;63(1):17–21.
10. Mor C, Abrahams VM. Immunology of implantation. *Immunol Allergy Clin N Am*. 2020;22(3):545–65.
11. Mor G, Cardenas I. The immune system in pregnancy :a unique complexity. *Am J Reprod Immunol*;2010(6):425–458.

12. Romero R, Espinoza J, Kusanovic JP, Gotsch F, Hassan S, Erez O. The preterm parturition syndrome BJOG. *Int J ObstetGynaecol.* 2006;113:17–42.
13. Bachmayer N, Sohlberg E, Sundstrom Y, Hamad RR, Berg L, Bremme K. Women with pre-eclampsia have an altered NKG2A and NKG2A receptor expression on peripheral blood natural killer cells. *Am J ReprodImmunol.* 2009;62(3):147–57.
14. Phoswa WN, Naicker T, Ramsuran V, Moodley J. Pre-eclampsia : the role of highly active antiretroviral therapy and immune markers. *Inflamm Res.* 2019;68(1):47–57.
15. Wan Y, Shang J, Graham R, Baric RS, Li F. Receptor recognition by the novel coronavirus from Wuhan : an analysis based on decade -long structural studies of SARS coronavirus. *J Virol.* 2020;94(7).
16. Hanna J, Goldmann-Wohl D, Hamani Y, Greenfield AI, Yaron CN, S. Decidual NK cells regulate key developmental processes at the human fetal -maternal interface. *Nat Med.* 2006;12(9):1065–74.
17. Chen Y, Liu Q, Guo D. Emerging coronaviruses :genome structure , replication , and pathogenesis. *J Med Virol.* 2020;92(4):418–441.
18. Shereen MA, Khan S, Kazmi A, Bashir N, Siddique R. COVID -19 infection :origin , transmission , and characteristics of human coronaviruses. *J Adv Res.* 2020;2:91–99.
19. Bourboulisej G, Netea MG, Rovina N, Akinosoglou K, Antoniadou A, Antonakos N. Complex immune dysregulation in COVID-19 patients with severe respiratory failure. *Cell Host Microbe.* 2020;27(6):992–1000.
20. Zheng M, Gao Y, Wang G, Song G, Liu S, Dun D. Functional exhaustion of antiviral lymphocytes in COVID -19 patients. *Cell Mol Immunol.* 2020;p. 1–3.
21. Ruan Q, Yang K, Wang W. Clinical Predictors of mortality due to COVID -19 based on analysis of data of 150 patients from Wuhan. *China Intensive Care Med.* 2020;p. 1–3.
22. Aghaeepour N, Ganio EA, Mcilwain D, Tsai SA, Tingle M, Gassen V. An immune clock of human pregnancy. *SciImmunol.* 2017;2(15).
23. Turco MY, Moffett A; 2019.
24. M, Zhang CI, Xiong J, Li C, X. The SARS-CoV -2 receptor ACE2 expression of maternal -fetal interface and fetal organs by single cell transcriptome study. *PLoS One.* 2020;15(4):230295–230295.
25. Mahase E, Covid. 19:most patients require mechanical ventilation in first 24 hours of critical care. *BMJ2020;368:m 1201doi.* 2020;.
26. Dashraath P, Jeslynw JL, Karen MX, I, Lim LM, Biswas LS, et al. COVID-19 )Pandemic and Pregnancy. *Coronavirus Disease.* 2019;(20):3033–3037.
27. Mcgonag LD, Sharif K, O’reagan A, Bridgewood C. The role of cytokines including interleukin-6 in COVID -19 induced pneumonia and macrophage activation syndrome -like disease. *Autoimmune Rev.* 2020;102537.
28. Morg C. The immune system in pregnancy : a unique complexity. *Am J ReprodImmunol.* 2010;63(6):425–458.
29. Wegmannt. Foetal protection against abortion: is it immunosuppression or immunomodulation ? *Ann Inst Pasteur /Immunol.* 1984;135(3):309–321.

**How to cite this article:** Kaur D.B. COVID 19 in Pregnancy: Immunology Savior or Culprit. *Journal of Medical Biomedical and Applied Sciences.* 2021;740–744. <https://doi.org/xx.xxx/xxx.xx>